

## Calibration of Radar Altimeters in the Corsica Area: Recent Improvements in view of Jason-1 mission



Y. Ménard (CNES), E. Jeansou (SILOGIC) and A. Orsoni (IGN) Centre National d'Etudes Spatiales, Toulouse, France

P. Exertier, P. Bonnefond, O. Laurain and F. Barlier Observatoire de la Côte d'Azur - CERGA, Grasse, France B. Haines (JPL), Pasadena, California, USA D.G. Kubitschek (CCAR), Boulder, Colorado, USA



## bonnefond@obs-azur.fr



Abstract
Since two years, we have developed a double geodetic site in Corsica (Western Mediterranean), in view of establishing a semi-permanent base for the purpose of absolute altimeter calibration experiments.


A first campaign has been carried out between 1996 - 1997 | Exertier et al.,
1998]. A fundamental good etic point has been established, thanks to the FILR,

1999). A "Humamora in gent point and service established, manual to the # 1.1-8.

Sty, and DOMS systems (Appricts, Morie Base, Hacio), in addition to a 1875, and DOMS systems (Appricts, Morie Base, Hacio), in addition to a 58, which is located at the Sentess Cap (40 km seath Ajaccio). Uncertainties on the good affittions at Sentess Auer Lead to a Calibration value of 5.6. a0.7-cm for T/P. In view of the Jason-I mission, and thanks to a GPS-away reprintent, which was conducted in May 1998, the Sentess six has been experiment, which was conducted in May 1998, the Sentess six has been experiment, which was conducted in May 1998, the Sentess six has been greatly improved.
The experiment is briefly described. First results of measurements of sea

levels from the tide gauges and buoys are given; comparisons are at the level of 0,7-1.5 cm rms, which is very encouraging for preparing the calibration of Jason-1 mission.

## General overview of the calibration site



# Brief History - Sept. 1999 to Feb. 1997: Calibration experiment using French Transportable Laser Ranging System at Ajaccio and the other as quages (one at Ajaccio and the other at Senetosa). - May 1998: Installation of two others tide gauges at Senetosa. Geodetic operations (GPS, leveling, see Tables and Cartering and Senetosa).

#### GPS Buovs Experiment

GPS BLOys Experiment


Calibration results from the first campaign (1996-97) have shown that the geold gradient in the Senetosa area should be very high. Then, it has been decided to improve this after by a state of the state o

Die gangen i GPS Broops	Gentelk politi
м,	6,
M <sub>2</sub>	8
	6,
SER CELGPS Box;	6,
ASS TEXTS OF BILL	g.
Parte regre GPS Berg	G.

Dis CPS	3,144	w) drg)	h (m)
G.	1.70325402617	40339034311722	85.5853
e.	8.798 del 13.772 26 8 3	4031201201210	52,712.6
6,	1.793 485 142 5 25 6 5	40334010310449	59.9319
o.	1.1146 7114 (202)	40249510 (5150)	31.5477

## **GPS Buoys experiment results**

#### Configuration in Corsica








Figure 2. GPS brows: Turbo-some (left) and SERCEL (right). Figure 3. Turbo-some GPS on G.

## 1- GPS Buoys Heights determination

#### instantaneous GPS Heights

Instantaneous GPS Heights

— GPS heights have been processed from RINEX files of all the receivers using Ashten's Software (PRISM-PNAV Vers. 24).

GPS orbits are from navigation message, inospheric orbits of the state of the participants (white-orbits of the participants (white-orbits or the state of the state orbits or the state of the state orbits of the state orbits or the state of the state orbits or the state orbits or the state of the state orbits or the state orbits or the state orbits or the state orbits orb

#### Filtered GPS Heights

— All GPS heights have been filtered using Vondrak filter [Vondrak, 1977], with a 60 second period. It means that all signals with period higher than 60 s are kept. An example of filtered data is given in Figure 4. — GPS heights from GOAII software (Turbo-rogue data) have



Figure 4, GPS heights from SERCEL data.
Red lines correspond to filtered heights using Vondrak filter

#### Comparisons

All GPS heights (filtered or not) have been compared for common sessions (different receivers).
 Comparisons of software solutions have also been performed.

Performed.

Results are given in Table 3. An example of differences between common sessions and 2 different receivers is presented in Figure 5.

Dab"		m is	891	801	HMCY	c som more all
Raw Brief	13451	461	4612 3.65	824 823	5.56 136	Tabo report SERCEL Tabo report SERCEL
Raw Named	1471	2379 3.88	22.36	1.4	437 1.62	Todo-copie - ASSITECH Todo-copie - ASSITECH
Raw Named	17632	-3164	435	1.34	2.4H 157	MERCEL MERCEL(DINF):
Raw Name	28033 28033	-4511	231	1.2	2.51	Timbo regare Technologia (GO A) (5) Timbo regare Technologia (GO A) (5)

Hat: Verdat wid 60 pried ha been applied for all de compasion.

\*Selvence SPACE from Darriet MECH. Noviguies Britishing.

\*Selvence Gipp Dark 1.



Figure 5. Differences between Turbo cogue and A shtech GPS heights on the 05/10/98 session.

#### 2- Tides correction for GPS Buoys Heights

— The GPS heights have been corrected from tides variations using mean tides variations computed with the three tide determination has been performed by 8. Haines with Turborogue data (GPS heights, filler, tides correction); differences are presented in Table 4.

— The Tide gauges time series are illustrated in Figure 6.

— An example, of this correction is given in Figure 7.





wen A. Omeni and B. Mainer relations, units are in continued

Raw	2818.6	-65.11	32.52	-1.56	438	Tinbe-regen (A. Orrent) - Terbe-regen (B. Haben)
Filters	2818.6	-11.24	8.68		2.42	Tinbe-regen (A. Orrent) - Terbe-regen (B. Haben)
The:	Vendrak wid	60 (proid)	Ceresian	fer sider free	P. Demefe	nd (Dured on die sverige of die 3 übe ganger dus) ;
	Alle Beson	- (GOA-II)	le revoluni	I Cerre o less	fie üler fie	im B. Rainer (Dured in ibe average of die 3 übe ganger dus) ;
Table 5	. Muse	fride gange	n beighen n	nd dute free	. 12 11112 1	(Fall) is \$511191 (Fall), from M., M., and M.,

	M,	M,	M,
Mean inte (CNES day) Mean beight (m)	1 ME 1 1 ME 1 733379	17(11)7511 47343	17441.37911 47.41.4
Reference gred: by 12 (05)	59 pt 2191		

Table 6.	$\textbf{6.}  \text{Corrections upplied to CP-Shedy has (compared from $M_{\star}, M_{\star}$ and $M_{\star}$), units are in continuous.}$				
	min	816	86.11	Hiller	common
75597 75597	1379 136	131	1.17	6.55 1.25	Mounths Sunded Deviation of man his

### 3- Comparisons with Tide gauges

— Differences between GPS heights and tide gauges heights have been computed at sech GPS data using linear interpolation between tide gauges location are given in Table 7. Figure 3 gives an exemple of such differences for M, tide gauge. Figure 9 shows QPS heights and M, tide gauge data for a QPS seession.

Table 7.	CPS Height - T					
Dab		min	B 16	8611	siller	commen
Raw Fired	3460	-1.33 -1.33	12.45	1.00	135	Alteriven:MI Alteriven:MI
Raw	3552 3552	-1.25 -1.38	11.54 3.78	1.0	251	Alteriven:M2 Alteriven:M2
Rev	3725	-6.81	7.0	1.5	161	All receives (M3

GPS Buoys - Tide gauge: M2

Figure 8. Differences between GPS and tide gauges heights at M, location. Red lines for filtered GPS heights.




Figure 9. GPS buoys and tide gauges heights at M. location on the 05/08/98.

#### 4- Geoid Height differences

Figure 10 illustrates the altitude differences. Where Q refers to Geold and E refers to Elipsoid.  $h_{\alpha} = h_{\alpha} + h_{\alpha}$ 

computing differences. Only  $a_0 = (a_0 + b_1)$  good height difference between points 1 and 2.  $(b_{11}, a_0) = (b_{11} + b_2)$  deduced from GPS heights.

• Oh,  $\alpha_{ij} = \{h_i, h_j\}$ ; deduced from GPS heights.

• Oh,  $\alpha_{ij} = \{h_i, h_j\}$ ; measured by leveling.

So Oh,  $\alpha_{ij} = Ah_{ij}$ ; measured by leveling.

Bor Oh,  $\alpha_{ij} = Ah_{ij}$ ; measured by leveling.


Boring the experiment geodelic points heights near each tide gauges have been measured by GPS and heights difference between M, and M, has been measured by leveling. Results are given in Table 9.

	GPSBm11	Tile pages"	Levin
Table 5. Computerners	reid beigde differmates ar s	g and form CPS banyo, ride	garger and leyding, min
Using mean of filtered dom, with a	de omecan.		
Sander Edition (m)	1318	1.0157	1319
Managini (mi	473588	67363	47.603
	me,	MA,	BM,



#### 5- Map of GPS Buovs Heights

Figure 11 illustrates the sea heights obtained with all GPS filtered heights corrected from tides. Interpolations for gridding have been limited to 2.8 km distance with a grid spacing of 50



8.	74	8.76	8.78	8.8	0	8.82
Binsize: Minimum Maximum	0.0250000 volue: 4 volue:	6,7880 47,4497	Longitude Int above CRS80	Stratout	of values: Mean: deviation:	75987 47,1987 0.161470
					-	
46.80	46.90	47.00	47.10	47.20	47.30	47.40
						21.5

#### Discussion

First, we want to emphasize how intensive the survey was and how rich the data set is. It is probably the first time that a site has been successfully surveyed so much with dual-redundant pelagic GPS, and complemented it with terrestrial GPS. 3 simultaneous tides and complemented it with three in a second of the property of the complement of the

#### GPS Buoys preliminary results

— <u>Comparisons of receivers/softwares;</u> differences are at the level of 1-2 cm rms with mean differences below 1 cm. The main signal appears to be blases by GPS session may be due to solving technique (lonospheric blas,

— <u>Comparisons with tide gauges (Table 7)</u>: GPS heights appear to be always higher than for tide gauges (-1.4 cm). The mean difference could be due in part to the presence of the radiome on one artenna (while the flucies) also have the presence of the radiome on one artenna (while the flucies) also have the presence of mean blases between (M, M), and M, have to be investigated (fide gauges blases).

— See heights have then be determined with a precision better than 1-2 cm rms, showing an along-track slope of 50 cm over 10 km and across-track gradient which can reach -5 cm between M2 and M3 (-3 km).

Use of the sea heights grid to link T/P measurements and tide gauges one in order to perform altimeter calibration.

- Densification of the GPS Buoys heights if necessary.

#### New celibration composign using FTLRS at Alaccio in 1999.

Exertier, P., P. Bonnefond, O. Laurein, F. Pierron, and F. Berlier, New French Transportable Laser Ranging System: positioning, collocation and altimeter cellibratic experiments results, submitted to Journal of Geodesy, 1998.

Blewitt, G., Carrier phase ambiguity resultion for the Global positioning system applied to geodetic baselines up to 2000 km, J. Geophys. Res. 94(B8), 10187-10293, 1989. Vondrak, J., Problem of smoothing observational data ii. Bull. Astron. Inst. Czech, vol. 28, pp. 84-89, 1977.

Webb, F. and J. Zumberge, An introduction to Gipsy/Oasis II, Precision sotware for the analysis of data from the global positioning system, JPL D-11088, 1995.