



# TOPEX/Poseidon MGDR Quality Assessment Report

Cycle 379

**28-12-2002 07-01-2003**

Prepared by :	C. Schgounn, CLS  G. Pontonnier, CLS  M. Ablain, CLS	
Accepted by :	J. Dorandeu, CLS	
Quality visa :	M. Destouesse, CLS	
Approved by :	N. Picot, CNES	



CENTRE NATIONAL D'ETUDES SPATIALES



**JPL**  
Jet Propulsion Laboratory  
California Institute of Technology

SALP-RP-P2-EX-21120-CLS379

Edition 01.0, June 2003

## **1 Introduction. Document overview**

The purpose of this document is to report the major features of the data quality from the Topex/Poseidon mission. The document is associated with data dissemination on a cycle by cycle basis.

The objectives of this document are :

- To provide a data quality assessment
- To provide users with necessary information for data processing
- To report any change likely to impact data quality at any level, from instrument status to software configuration
- To present the major useful results for the current cycle

It is divided into the following topics:

- [\*\*Cycle overview\*\*](#)
- [\*\*CALVAL main results\*\*](#)

## 2 Cycle overview


### 2.1 Cycle quality and performances

Data quality for this cycle appears to be nominal. For this cycle, the crossover standard deviation is 6.42 cm rms, and the standard deviation of Sea Level Anomalies (SLA) relative to a Mean Sea Surface is 9.66 cm.

### 2.2 Warnings and recommendations

- Missing measurements :  
Passes 3, 14, 105, 142 and 168 are missing due to small size. All the passes are located above the Indian Ocean.
- Tape recorder failures :  
There is a lot of data gaps due to tape recorder anomalies, especially in the Indian Ocean.
- Editing measurements (a) :  
Problems in the interpolation of the TMR parameters occur when there are missing measurements (tape recorder failures). As a result 4.46% of the measurements are removed by the TMR correction criterion.
- Editing measurements (b) :  
A new criterion has been added to the editing procedure since cycle 376 (See [Editing](#)). The measurements removed by this criterion for the current cycle are plotted on the following figure.

Edited parameter : Radiometer and ECMWF wet tropospheric differences  
T/P Cycle 379 (28/12/2002 / 07/01/2003)




### 3 CALVAL main results

This section presents results that illustrate data quality during this cycle. These verification products are produced operationally so that they allow systematic monitoring of the main relevant parameters.

#### 3.1 Missing measurements

698431 altimeter measurements are present, and 96081 are missing.

The map below shows all the available measurements for this cycle and illustrates the tape recorder problems. The latter figure shows missing 1Hz measurements in the GDRs, with respect to a 1 Hz sampling of a nominal repeat track.




### 3.2 M-GDR quality flags


The following table indicates the percentage of measurements for which those flags are set.

Name	Description	% bad
Geo_Bad_1	altimeter land flag	26.33
Geo_Bad_1	ice flag	4.47
Geo_Bad_1	radiometer land flag	27.99
Alt_Bad_1	conditions 1 altimeter	4.23
Alt_Bad_2	conditions 2 altimeter	4.12
Geo_Bad_2	rain (liquid water in excess)	6.77
Geo_Bad_2	less than 4 points for CSR3.0 tide calculation	0.36
Geo_Bad_2	less than 4 points for FES95.2.1 tide calculation	2.44
TOPEX	TOPEX not valid	0.00
TMR	TMR not valid	0.00
TMR_Bad	Brightness temperatures not valid	4.40
DORIS	DORIS not valid	0.00


### 3.3 M-GDR parameter plots




Cycle 379 – GDR\_M – TOPEX – All passes –







Cycle 379 – GDR\_M – TOPEX – All passes –




Cycle 379 – GDR\_M – TOPEX – All passes –









### 3.4 Editing

The following table gives for each tested parameter, minimum and maximum thresholds, the number and the percentage of points removed.


As a comparison, the mean percentage over one year (1997) is also given.

There are problems in the interpolation of the TMR parameters since cycle 371 when there are missing measurements (tape recorder failures). These bad measurements are removed by the TMR correction criterion but some of them have been kept. Thus a new criterion has been added to the editing procedure since the cycle 376 to remove all the measurements where the absolute value of the difference between the TMR correction and the ECMWF model wet tropospheric correction is greater than 20 cm.


Parameters	Min Thres.	Max Thres.	Unit	Mean removed in 1997	% removed
Sea surface height	-130.000	100.000	m	1.37	0.45
Number of 20/10Hz valid points Poseidon/TOPEX	5.000	-		1.37	0.60
Std. deviation of range	0.000	0.100	m	1.85	1.39
Off nadir angle from waveform	0.000	0.400	deg	1.36	3.55
Dry tropospheric correction	-2.500	-1.900	m	0.00	0.00
Invert barometer correction	-2.000	2.000	m	0.00	0.00
TMR wet tropospheric correction	-0.500	-0.001	m	0.34	4.46
Ionospheric correction (Poseidon:Doris, TOPEX:Dual)	-0.400	0.040	m	0.00	0.00
Significant wave height	0.000	11.000	m	1.46	0.25
Sea state Bias	-0.500	0.000	m	1.39	0.40
Backscatter coefficient	7.000	30.000	dB	1.44	0.39
Ocean tide height	-5.000	5.000	m	0.01	0.51
Earth tide	-1.000	1.000	m	0.00	0.00
Pole tide	-15.000	15.000	m	0.00	0.00
TMR and ECMWF tropospheric differences	-0.200	0.200	m	NaN	0.55
Spline fitting					0.01

The following three maps are complementary: they show respectively the removed, the selected measurements and the percentage of selected measurements in the editing procedure.


Valid data  
TOPEX/Poseidon Cycle 379 (28/12/2002 / 07/01/2003)



Edited measurements  
TOPEX Cycle 379 (28/12/2002 / 07/01/2003)




Percentage of valid data relative to the nominal pass  
TOPEX/Poseidon Cycle 379 (28/12/2002 / 07/01/2003)




### 3.5 Ionospheric correction


TOPEX/Poseidon Cycle 379 ( 28/12/2002 – 07/01/2003 ), ascending passes




# TOPEX/Poseidon Cycle 379 ( 28/12/2002 – 07/01/2003 ), descending passes




TOPEX/Poseidon Cycle 379 ( 28/12/2002 – 07/01/2003 ), ascending passes




# TOPEX/Poseidon Cycle 379 ( 28/12/2002 – 07/01/2003 ), descending passes




TOPEX/Poseidon Cycle 379 ( 28/12/2002 – 07/01/2003 ), ascending passes




# TOPEX/Poseidon Cycle 379 ( 28/12/2002 – 07/01/2003 ), descending passes



### 3.6 Wet tropospheric corection



### 3.7 Crossover statistics



## TPTP – CROSSOVER STATISTICS

Without orbit error correction

**SSH = Corrected sea surface height**

### **RAPPEL DES SELECTIONS**

Type de points de croisement: TPTP

Zone geographique (deg): -90 / 90 , 0 / 360

Seuil sur les ecart s d'analyse 0.00 (moy)  
30.00 (seuil)

Selection(s) sur les champs :

CL Arc 1 :=INTERP\_SPLN

CL Arc 2 :=INTERP\_SPLN

Seuil Min +: 0.0000000

Seuil Max : 0.0000000

Selection(s) sur les ecart s :

Aucune

### **RESULTATS STATISTIQUES**

Valeur minimale : -29.6200

Valeur maximale : 29.7100

Difference Max – Min: 59.3300

Nombre de points lus: 5087


Nombre de points selectionnes: 4968

Moyenne : 0.0945350

Ecart-type : 6.45021

Moyenne Quadratique : 6.45090

**CLS Space Oceanography Division**



## TPTP – CROSSOVER STATISTICS

With orbit error correction

**SSH = Corrected sea surface height**

### **RAPPEL DES SELECTIONS**

Type de points de croisement: TPTP

Zone geographique (deg): -90 / 90 , 0 / 360

Seuil sur les ecart s d'analyse 0.00 (moy)  
30.00 (seuil)

Selection(s) sur les champs :

CL Arc 1 :=INTERP\_SPLN

CL Arc 2 :=INTERP\_SPLN

Seuil Min +: 0.0000000

Seuil Max : 0.0000000

Selection(s) sur les ecart s :

Aucune

### **RESULTATS STATISTIQUES**

Valeur minimale : -29.5100

Valeur maximale : 29.0600

Difference Max – Min: 58.5700

Nombre de points lus: 5087


Nombre de points selectionnes: 4954

Moyenne : 0.0577170

Ecart-type : 5.69252

Moyenne Quadratique : 5.69282

**CLS Space Oceanography Division**



# TPTP – CROSSOVER STATISTICS

SSH, BATHY < -1000 m, VAR\_OCE < 20 cm, LAT [-50°, +50]

**SSH = Corrected sea surface height before orbit error**

## **RAPPEL DES SELECTIONS**

Type de points de croisement: TPTP  
 Zone géographique (deg): -50 / 50 , 0 / 360  
 Seuil sur les écarts d'analyse : aucun

Selection(s) sur les champs :


CL Arc 1 :=BATHY  
 CL Arc 2 :=BATHY  
 Seuil Min : aucun  
 Seuil Max : -1000000.00  
 CL Arc 1 :=VAR\_OCE  
 CL Arc 2 :=VAR\_OCE  
 Seuil Min : aucun  
 Seuil Max : 20.000000  
 [...]

Selection(s) sur les écarts :  
 Aucune

## **RESULTATS STATISTIQUES**

Valeur minimale : -49.2600  
 Valeur maximale : 38.7200  
 Différence Max – Min: 87.9800  
 Nombre de points lus: 2762  
 Nombre de points sélectionnés: 2521  
 Moyenne : 0.0671720  
 Ecart-type : 5.88111  
 Moyenne Quadratique : 5.88150


**CLS Space Oceanography Division**




## 3.8 SSH variability

### 3.8.1 Sea Level Anomaly


TOPEX/Poseidon, cycle 379  
Period : 28/12/2002 – 07/01/2003



TOPEX/Poseidon, cycle 379  
Period : 28/12/2002 – 07/01/2003




TOPEX/Poseidon, cycle 379  
Period : 28/12/2002 – 07/01/2003




### 3.8.2 Comparison to a precise Mean Sea Surface

The CLS (2001) MSS model is used as a reference to compute SLA. The two following maps respectively show the map of Topex SLA relative to the MSS and differences higher than a 30 cm threshold (after centering the data). The latter figure shows that higher differences are located in high ocean variability areas, as expected.



### 3.9 Dynamic topography


TOPEX/Poseidon, cycle 379  
Period : 28/12/2002 – 07/01/2003



### 3.10 Wind and wave maps

These two figures show wind and wave estimations derived from 10 days of altimeter measurements.

TOPEX/Poseidon, cycle 379  
Period : 28/12/2002 – 07/01/2003



TOPEX/Poseidon, cycle 379  
Period : 28/12/2002 – 07/01/2003

