



TOPEX/Poseidon MGDR Quality Assessment Report

Cycle 381

17-01-2003 27-01-2003

Prepared by :	C. Schgounn, CLS G. Pontonnier, CLS M. Ablain, CLS	
Accepted by :	J. Dorandeu, CLS	
Quality visa :	M. Destouesse, CLS	
Approved by :	N. Picot, CNES	



CENTRE NATIONAL D'ETUDES SPATIALES



JPL
Jet Propulsion Laboratory
California Institute of Technology

SALP-RP-P2-EX-21120-CLS381

Edition 01.0, June 2003

1 Introduction. Document overview

The purpose of this document is to report the major features of the data quality from the Topex/Poseidon mission. The document is associated with data dissemination on a cycle by cycle basis.

The objectives of this document are :

- To provide a data quality assessment
- To provide users with necessary information for data processing
- To report any change likely to impact data quality at any level, from instrument status to software configuration
- To present the major useful results for the current cycle

It is divided into the following topics:

- Cycle overview**
- CALVAL main results**

2 Cycle overview

2.1 Cycle quality and performances

Data quality for this cycle appears to be nominal. For this cycle, the crossover standard deviation is 6.33 cm rms, and the standard deviation of Sea Level Anomalies (SLA) relative to a Mean Sea Surface is 9.47 cm.

2.2 Warnings and recommendations

- Tape recorder failures :


There are a lot of data gaps due to tape recorder anomalies, especially in the Indian Ocean.

- Editing measurements (a) :

Problems in the interpolation of the TMR parameters occur when there are missing measurements (tape recorder failures). As a result 4.71% of the measurements are removed by the TMR correction criterion.

- Editing measurements (b) :

A new criterion has been added to the editing procedure since cycle 376 (See [Editing](#)). The measurements removed by this criterion for the current cycle are plotted on the following figure.




3 CALVAL main results

This section presents results that illustrate data quality during this cycle. These verification products are produced operationally so that they allow systematic monitoring of the main relevant parameters.

3.1 Missing measurements

701240 altimeter measurements are present, and 93288 are missing.

The map below shows all the available measurements for this cycle and illustrates the tape recorder problems. The latter figure shows missing 1Hz measurements in the GDRs, with respect to a 1 Hz sampling of a nominal repeat track.





3.2 M-GDR quality flags

The following table indicates the percentage of measurements for which those flags are set.


Name	Description	% bad
Geo_Bad_1	altimeter land flag	27.19
Geo_Bad_1	ice flag	3.51
Geo_Bad_1	radiometer land flag	28.40
Alt_Bad_1	conditions 1 altimeter	4.22
Alt_Bad_2	conditions 2 altimeter	4.11
Geo_Bad_2	rain (liquid water in excess)	6.00
Geo_Bad_2	less than 4 points for CSR3.0 tide calculation	0.35
Geo_Bad_2	less than 4 points for FES95.2.1 tide calculation	2.46
TOPEX	TOPEX not valid	0.00
TMR	TMR not valid	0.00
TMR_Bad	Brightness temperatures not valid	5.88
DORIS	DORIS not valid	0.00

3.3 M-GDR parameter plots










Cycle 381 – GDR_M – TOPEX – All passes –




Cycle 381 – GDR_M – TOPEX – All passes –









3.4 Editing

The following table gives for each tested parameter, minimum and maximum thresholds, the number and the percentage of points removed.


As a comparison, the mean percentage over one year (1997) is also given.

There are problems in the interpolation of the TMR parameters since cycle 371 when there are missing measurements (tape recorder failures). These bad measurements are removed by the TMR correction criterion but some of them have been kept. Thus a new criterion has been added in the editing procedure since cycle 376 to remove all the mesurements for the difference between the TMR and the ECMWF tropospheric corrections is greater than 20 cm. Even after applying this editing procedure, some anomalous TMR correction values remain (See [wet tropo correction section](#)).


Parameters	Min Thres.	Max Thres.	Unit	Mean removed in 1997	% removed
Sea surface height	-130.000	100.000	m	1.37	1.34
Number of 20/10Hz valid points Poseidon/TOPEX	5.000	-		1.37	1.65
Std. deviation of range	0.000	0.100	m	1.85	2.49
Off nadir angle from waveform	0.000	0.400	deg	1.36	4.72
Dry tropospheric correction	-2.500	-1.900	m	0.00	0.00
Invert barometer correction	-2.000	2.000	m	0.00	0.00
TMR wet tropospheric correction	-0.500	-0.001	m	0.34	4.71
Ionospheric correction (Poseidon:Doris, TOPEX:Dual)	-0.400	0.040	m	0.00	0.00
Significant wave height	0.000	11.000	m	1.46	0.75
Sea state Bias	-0.500	0.000	m	1.39	1.19
Backscatter coefficient	7.000	30.000	dB	1.44	1.09
Ocean tide height	-5.000	5.000	m	0.01	1.63
Earth tide	-1.000	1.000	m	0.00	0.00
Pole tide	-15.000	15.000	m	0.00	0.00
TMR and ECMWF tropospheric differences	-0.200	0.200	m	NaN	1.61
Spline fitting					0.01

The following three maps are complementary: they show respectively the removed, the selected measurements and the percentage of selected measurements in the editing procedure.


Valid data
TOPEX/Poseidon Cycle 381 (17/01/2003 / 27/01/2003)



Edited measurements
TOPEX Cycle 381 (17/01/2003 / 27/01/2003)




Percentage of valid data relative to the nominal pass
TOPEX/Poseidon Cycle 381 (17/01/2003 / 27/01/2003)




3.5 Ionospheric correction


TOPEX/Poseidon Cycle 381 (17/01/2003 – 27/01/2003), ascending passes




TOPEX/Poseidon Cycle 381 (17/01/2003 – 27/01/2003), descending passes




TOPEX/Poseidon Cycle 381 (17/01/2003 – 27/01/2003), ascending passes




TOPEX/Poseidon Cycle 381 (17/01/2003 – 27/01/2003), descending passes




TOPEX/Poseidon Cycle 381 (17/01/2003 – 27/01/2003), ascending passes




TOPEX/Poseidon Cycle 381 (17/01/2003 – 27/01/2003), descending passes





3.6 Wet tropospheric corection



3.7 Crossover statistics





TPTP – CROSSOVER STATISTICS

SSH, BATHY < -1000 m, VAR_OCE < 20 cm, LAT [-50°, +50]

SSH = Corrected sea surface height before orbit error

RAPPEL DES SELECTIONS

Type de points de croisement: TPTP
 Zone géographique (deg): -50 / 50 , 0 / 360
 Seuil sur les écarts d'analyse : aucun

Selection(s) sur les champs :

CL Arc 1 :=BATHY
 CL Arc 2 :=BATHY
 Seuil Min : aucun
 Seuil Max : -100000.00
 CL Arc 1 :=VAR_OCE
 CL Arc 2 :=VAR_OCE
 Seuil Min : aucun
 Seuil Max : 20.000000
 [...]


Selection(s) sur les écarts :

Aucune

RESULTATS STATISTIQUES

Valeur minimale : -31.1500
 Valeur maximale : 38.9300
 Différence Max – Min: 70.0800
 Nombre de points lus: 2673
 Nombre de points sélectionnés: 2431
 Moyenne : 0.606602
 Ecart-type : 5.77358
 Moyenne Quadratique : 5.80536


CLS Space Oceanography Division




3.8 SSH variability

3.8.1 Sea Level Anomaly


TOPEX/Poseidon, cycle 381
Period : 17/01/2003 – 27/01/2003



TOPEX/Poseidon, cycle 381
Period : 17/01/2003 – 27/01/2003




TOPEX/Poseidon, cycle 381
Period : 17/01/2003 – 27/01/2003




3.8.2 Comparison to a precise Mean Sea Surface

The CLS (2001) MSS model is used as a reference to compute SLA. The two following maps respectively show the map of Topex SLA relative to the MSS and differences higher than a 30 cm threshold (after centering the data). The latter figure shows that higher differences are located in high ocean variability areas, as expected.



3.9 Dynamic topography


TOPEX/Poseidon, cycle 381
Period : 17/01/2003 – 27/01/2003



3.10 Wind and wave maps

These two figures show wind and wave estimations derived from 10 days of altimeter measurements.

TOPEX/Poseidon, cycle 381
Period : 17/01/2003 – 27/01/2003



TOPEX/Poseidon, cycle 381
Period : 17/01/2003 – 27/01/2003

