



TOPEX/Poseidon MGRD Quality Assessment Report

Cycle 396

14-06-2003 24-06-2003

Prepared by :	C. Schgounn, CLS G. Pontonnier, CLS M. Ablain, CLS	
Accepted by :	J. Dorandeu, CLS	
Quality visa :	M. Destouesse, CLS	
Approved by :	N. Picot, CNES	



SALP-RP-P2-EX-21120-CLS396

Edition 01.0, August 2003

1 Introduction. Document overview

The purpose of this document is to report the major features of the data quality from the Topex/Poseidon mission. The document is associated with data dissemination on a cycle by cycle basis.

The objectives of this document are :

- To provide a data quality assessment
- To provide users with necessary information for data processing
- To report any change likely to impact data quality at any level, from instrument status to software configuration
- To present the major useful results for the current cycle

It is divided into the following topics:

[Cycle overview](#)

[CALVAL main results](#)

2 Cycle overview


2.1 Cycle quality and performances

Data quality for this cycle appears to be nominal. For this cycle, the crossover standard deviation is 6.89 cm rms, and the standard deviation of Sea Level Anomalies (SLA) relative to a Mean Sea Surface is 11.54 cm.

2.2 Warnings and recommendations

- Missing measurements :
There is a lot of data gaps due to tape recorder anomalies, especially in the Indian Ocean and in the Pacific Ocean below the South America coasts.
- DORIS incident :
The DORIS instrument experienced an incident on 16, June 2003. Therefore there is no DORIS data available from 17/06/2003 to 19/06/2003. Only bent ionospheric correction is available during this period, that is from pass 57 to 133.
- Measurements edited by the TMR parameters :
The following anomalies are explained by the problems in the interpolation of the TMR parameters due to tape recorder failures :
 - 3.23% of the measurements are removed by the TMR correction criterion (see the following figure).
 - Some passes have radiometer earth flag set to valid over earth. But these measurements are not kept and edited by the altimetric parameters.

Edited parameter : Radiometer wet tropospheric correction
T/P Cycle 396 (14/06/2003 / 24/06/2003)



3 CALVAL main results


This section presents results that illustrate data quality during this cycle. These verification products are produced operationally so that they allow systematic monitoring of the main relevant parameters.

3.1 Missing measurements


691858 altimeter measurements are present, and 102703 are missing.

The map below shows all the available measurements for this cycle and illustrates the tape recorder problems. The latter figure shows missing 1Hz measurements in the GDRs, with respect to a 1 Hz sampling of a nominal repeat track.

Available measurements
TOPEX Cycle 396 (14/06/2003 / 24/06/2003)



Missing measurements
TOPEX/Poseidon Cycle 396 (14/06/2003 / 24/06/2003)




3.2 M-GDR quality flags

The following table indicates the percentage of measurements for which those flags are set.


Name	Description	% bad
Geo_Bad_1	altimeter land flag	25.89
Geo_Bad_1	ice flag	7.38
Geo_Bad_1	radiometer land flag	27.83
Alt_Bad_1	conditions 1 altimeter	5.03
Alt_Bad_2	conditions 2 altimeter	4.90
Geo_Bad_2	rain (liquid water in excess)	5.40
Geo_Bad_2	less than 4 points for CSR3.0 tide calculation	0.40
Geo_Bad_2	less than 4 points for FES95.2.1 tide calculation	2.81
TOPEX	TOPEX not valid	0.00
TMR	TMR not valid	0.00
TMR_Bad	Brightness temperatures not valid	5.18
DORIS	DORIS not valid	0.00

3.3 M-GDR parameter plots


Cycle 396 – GDR_M – TOPEX – All passes –




Cycle 396 – GDR_M – TOPEX – All passes –










Cycle 396 – GDR_M – TOPEX – All passes –




Cycle 396 – GDR_M – TOPEX – All passes –









3.4 Editing

The following table gives for each tested parameter, minimum and maximum thresholds, the number and the percentage of points removed.

As a comparison, the mean percentage over one year (1997) is also given.


There are problems in the interpolation of the TMR parameters since cycle 371 when there are missing measurements (tape recorder failures). These bad measurements are removed by the TMR correction criterion but some of them have been kept. Thus a new criterion has been added to the editing procedure since the cycle 376 to remove all the measurements where the absolute value of the difference between the TMR correction and the ECMWF model wet tropospheric correction is greater than 20 cm.

The number and percentage of points removed by each criterion is given on the following table. Note that these statistics are obtained with measurements already edited for radiometer land flag (27.83 % of points removed) and ice flag (7.38 % of points removed).


Parameters	Min Thres.	Max Thres.	Unit	Mean removed in 1997	% removed
Sea surface height	-130.000	100.000	m	1.37	0.56
Number of 20/10Hz valid points Poseidon/TOPEX	5.000	-		1.37	0.74
Std. deviation of range	0.000	0.100	m	1.85	1.59
Off nadir angle from waveform	0.000	0.400	deg	1.36	3.97
Dry tropospheric correction	-2.500	-1.900	m	0.00	0.00
Invert barometer correction	-2.000	2.000	m	0.00	0.00
TMR wet tropospheric correction	-0.500	-0.001	m	0.34	3.23
Ionospheric correction (Poseidon:Doris, TOPEX: Dual)	-0.400	0.040	m	0.00	0.00
Significant wave height	0.000	11.000	m	1.46	0.31
Sea state Bias	-0.500	0.000	m	1.39	0.49
Backscatter coefficient	7.000	30.000	dB	1.44	0.43
Ocean tide height	-5.000	5.000	m	0.01	0.66
Earth tide	-1.000	1.000	m	0.00	0.00
Pole tide	-15.000	15.000	m	0.00	0.00
TMR and ECMWF tropospheric differences	-0.200	0.200	m	NaN	0.67
Spline fitting					0.02

The following three maps are complementary: they show respectively the removed, the selected measurements and the percentage of selected measurements in the editing procedure.


Valid data
TOPEX/Poseidon Cycle 396 (14/06/2003 / 24/06/2003)



Edited measurements
TOPEX Cycle 396 (14/06/2003 / 24/06/2003)




Percentage of valid data relative to the nominal pass
TOPEX/Poseidon Cycle 396 (14/06/2003 / 24/06/2003)




3.5 Ionospheric correction

TOPEX/Poseidon Cycle 396 (14/06/2003 – 24/06/2003), ascending passes



minimum number of samples per box 1
maximum number of samples per box 5153



Statistics Y-X

mean = -0.01288
rms = 0.01475
std = 0.00719


Order 1 fit polynomial

$y = a x + b$
a = 0.54969621
b = -0.00133827


Legend

— Order 1 fit polynomial
— Bisectrix

TOPEX/Poseidon Cycle 396 (14/06/2003 – 24/06/2003), descending passes



minimum number of samples per box 1
 maximum number of samples per box 1005



Statistics Y-X

mean = -0.01442
 rms = 0.01926
 std = 0.01277


Order 1 fit polynom

$y = a x + b$
 $a = 0.78769630$
 $b = -0.00156694$


Legend

— Order 1 fit polynom
 - - - Bisectrix

TOPEX/Poseidon Cycle 396 (14/06/2003 – 24/06/2003), ascending passes



minimum number of samples per box 1
 maximum number of samples per box 4428



Statistics Y-X

mean = -0.01412
 rms = 0.01708
 std = 0.00962


Order 1 fit polynom

$y = a x + b$
 $a = 0.38497892$
 $b = 0.00180641$


Legend

— Order 1 fit polynom
 — Bisectrix


TOPEX/Poseidon Cycle 396 (14/06/2003 – 24/06/2003), descending passes



TOPEX/Poseidon Cycle 396 (14/06/2003 – 24/06/2003), ascending passes



minimum number of samples per box 5
 maximum number of samples per box 11258



Statistics Y-X

mean = -0.00090
 rms = 0.00434
 std = 0.00425


Order 1 fit polynom

$y = a x + b$
 $a = 0.77404803$
 $b = 0.00198642$


Legend

— Order 1 fit polynom
 — Bisectrix


TOPEX/Poseidon Cycle 396 (14/06/2003 – 24/06/2003), descending passes



3.6 Wet tropospheric correction



3.7 Crossover statistics



TPTP – CROSSOVER STATISTICS

Without orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS


Type de points de croisement: TPTP
 Zone géographique (deg): -90 / 90 , 0 / 360
 Seuil sur les écarts d'analyse 0.00 (moy)
 30.00 (seuil)
 Selection(s) sur les champs :
 CL Arc 1 :=INTERP_SPLN
 CL Arc 2 :=INTERP_SPLN
 Seuil Min +: 0.0000000
 Seuil Max : 0.0000000

Selection(s) sur les écarts :
 Aucune

RESULTATS STATISTIQUES

Valeur minimale : -29.7700
 Valeur maximale : 28.6000
 Différence Max – Min: 58.3700
 Nombre de points lus: 4758
 Nombre de points sélectionnés: 4646
 Moyenne : 0.850839
 Ecart-type : 7.02577
 Moyenne Quadratique : 7.07711

CLS Space Oceanography Division



TPTP – CROSSOVER STATISTICS

With orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS


Type de points de croisement: TPTP
 Zone géographique (deg): -90 / 90 , 0 / 360
 Seuil sur les écarts d'analyse 0.00 (moy)
 30.00 (seuil)
 Selection(s) sur les champs :
 CL Arc 1 :=INTERP_SPLN
 CL Arc 2 :=INTERP_SPLN
 Seuil Min +: 0.0000000
 Seuil Max : 0.0000000

Selection(s) sur les écarts :
 Aucune

RESULTATS STATISTIQUES

Valeur minimale : -29.7000
 Valeur maximale : 29.0900
 Différence Max – Min: 58.7900
 Nombre de points lus: 4758
 Nombre de points sélectionnés: 4630
 Moyenne : 0.0871730
 Ecart-type : 5.91555
 Moyenne Quadratique : 5.91620

CLS Space Oceanography Division



TPTP – CROSSOVER STATISTICS
SSH, BATHY < -1000 m, VAR_OCE < 20 cm, LAT [-50°, +50]
SSH = Corrected sea surface height before orbit error

RAPPEL DES SELECTIONS


Type de points de croisement: TPTP
 Zone géographique (deg): -50 / 50 , 0 / 360
 Seuil sur les écarts d'analyse : aucun
 Selection(s) sur les champs :
 CL Arc 1 : =BATHY
 CL Arc 2 : =BATHY
 Seuil Min : aucun
 Seuil Max : -100000.00
 CL Arc 1 : =VAR_OCE
 CL Arc 2 : =VAR_OCE
 Seuil Min : aucun
 Seuil Max : 20.000000
 [...]

 Selection(s) sur les écarts :
 Aucune

RESULTATS STATISTIQUES

Valeur minimale : -31.6800
 Valeur maximale : 46.2700
 Différence Max – Min: 77.9500
 Nombre de points lus: 2659
 Nombre de points selectionnes: 2430
 Moyenne : 0.621609
 Ecart-type : 5.91648
 Moyenne Quadratique : 5.94905


CLS Space Oceanography Division




3.8 SSH variability

3.8.1 Sea Level Anomaly


TOPEX/Poseidon, cycle 396
Period : 14/06/2003 – 24/06/2003



TOPEX/Poseidon, cycle 396
Period : 14/06/2003 – 24/06/2003




TOPEX/Poseidon, cycle 396
Period : 14/06/2003 – 24/06/2003




3.8.2 Comparison to a precise Mean Sea Surface

The CLS (2001) MSS model is used as a reference to compute SLA. The two following maps respectively show the map of Topex SLA relative to the MSS and differences higher than a 30 cm threshold (after centering the data). The latter figure shows that higher differences are located in high ocean variability areas, as expected.

TOPEX/Poseidon, cycle 396
Period : 14/06/2003 – 24/06/2003




(SSH - MSS) differences greater than 0.3 m
TOPEX/Poseidon Cycle 396 (14/06/2003 / 24/06/2003)



3.9 Dynamic topography


TOPEX/Poseidon, cycle 396
Period : 14/06/2003 – 24/06/2003



3.10 Wind and wave maps

These two figures show wind and wave estimations derived from 10 days of altimeter measurements.

TOPEX/Poseidon, cycle 396
Period : 14/06/2003 – 24/06/2003



TOPEX/Poseidon, cycle 396
Period : 14/06/2003 – 24/06/2003

