



TOPEX/Poseidon MGRD Quality Assessment Report

Cycle 408

11-10-2003 21-10-2003

Prepared by :	C. Schgounn, CLS G. Pontonnier, CLS M. Ablain, CLS	
Accepted by :	J. Dorandeu, CLS	
Quality visa :	M. Destouesse, CLS	
Approved by :	N. Picot, CNES	



1 Introduction. Document overview

The purpose of this document is to report the major features of the data quality from the Topex/Poseidon mission. The document is associated with data dissemination on a cycle by cycle basis.

The objectives of this document are :

- To provide a data quality assessment
- To provide users with necessary information for data processing
- To report any change likely to impact data quality at any level, from instrument status to software configuration
- To present the major useful results for the current cycle

It is divided into the following topics:

[Cycle overview](#)

[CALVAL main results](#)

2 Cycle overview

2.1 Cycle quality and performances

Data quality for this cycle appears to be nominal. For this cycle, the crossover standard deviation is 6.47 cm rms, and the standard deviation of Sea Level Anomalies (SLA) relative to a Mean Sea Surface is 9.67 cm.


2.2 Warnings and recommendations

- Missing measurements :
 - There is a lot of data gaps due to tape recorder anomalies, especially in the Indian Ocean, in the South Pacific Ocean close to the South and Central America coasts and below the Groenland coasts.
- Measurements edited by the TMR parameters :

The following anomalies are explained by the problems in the interpolation of the TMR parameters due to tape recorder failures :

 - 1.91% of the measurements are removed by the TMR correction criterion (see the following figure).
 - Some measurements have radiometer earth flag set to valid over earth. A new criterion has been added to the editing procedure to remove all these measurements (see [Editing](#)) .

Edited parameter : Radiometer wet tropospheric correction
T/P Cycle 408 (11/10/2003 / 21/10/2003)



3 CALVAL main results


This section presents results that illustrate data quality during this cycle. These verification products are produced operationally so that they allow systematic monitoring of the main relevant parameters.

3.1 Missing measurements


696610 altimeter measurements are present, and 97894 are missing.

The map below shows all the available measurements for this cycle and illustrates the tape recorder problems. The latter figure shows missing 1Hz measurements in the GDRs, with respect to a 1 Hz sampling of a nominal repeat track.

Available measurements
TOPEX Cycle 408 (11/10/2003 / 21/10/2003)



Missing measurements
TOPEX/Poseidon Cycle 408 (11/10/2003 / 21/10/2003)




3.2 M-GDR quality flags

The following table indicates the percentage of measurements for which those flags are set.






Name	Description	% bad
Geo_Bad_1	altimeter land flag	26.34
Geo_Bad_1	ice flag	8.35
Geo_Bad_1	radiometer land flag	28.20
Alt_Bad_1	conditions 1 altimeter	5.05
Alt_Bad_2	conditions 2 altimeter	4.92
Geo_Bad_2	rain (liquid water in excess)	5.42
Geo_Bad_2	less than 4 points for CSR3.0 tide calculation	0.44
Geo_Bad_2	less than 4 points for FES95.2.1 tide calculation	3.09
TOPEX	TOPEX not valid	0.00
TMR	TMR not valid	0.00
TMR_Bad	Brightness temperatures not valid	5.60
DORIS	DORIS not valid	0.00

3.3 M-GDR parameter plots


Cycle 408 – GDR_M – TOPEX – All passes –




Cycle 408 - GDR_M - TOPEX - All passes -









Cycle 408 – GDR_M – TOPEX – All passes –




Cycle 408 – GDR_M – TOPEX – All passes –









3.4 Editing

The following table gives for each tested parameter, minimum and maximum thresholds, the number and the percentage of points removed. As a comparison, the mean percentage over one year (1997) is also given.

There are problems in the interpolation of the TMR parameters since cycle 371 when there are missing measurements (tape recorder failures). These bad measurements are removed by the TMR correction criterion but some of them have been kept. Thus a new criterion has been added to the editing procedure since the cycle 376 to remove all the measurements where the absolute value of the difference between the TMR correction and the ECMWF model wet tropospheric correction is greater than 20 cm.


Probably due to the interpolation problem with the TMR, some measurements have radiometer land flag unset over land. This has no impact on the valid data because these measurements have been edited by the altimetric parameter criteria. Nevertheless, this anomaly leads to wrong statistics of the edited measurements. Therefore a new criterion has been added in the editing procedure to remove all the measurements for which the radiometer land flag is set to ocean and the altimeter land flag is set to land.

The number and percentage of points removed by each criterion is given on the following table. Note that these statistics are obtained with measurements already edited for radiometer land flag (28.20 % of points removed) and ice flag (8.35 % of points removed).


Parameters	Min Thres.	Max Thres.	Unit	Mean % removed in 1997	% removed
Sea surface height	-130.000	100.000	m	1.37	0.16
Number of 20/10Hz valid points Poseidon/TOPEX	5.000	-		1.37	0.24
Std. deviation of range	0.000	0.100	m	1.85	1.04
Off nadir angle from waveform	0.000	0.400	deg	1.36	3.42
Dry tropospheric correction	-2.500	-1.900	m	0.00	0.00
Invert barometer correction	-2.000	2.000	m	0.00	0.00
TMR wet tropospheric correction	-0.500	-0.001	m	0.34	1.91
Ionospheric correction (Poseidon:Doris, TOPEX:Dual)	-0.400	0.040	m	0.00	0.00
Significant wave height	0.000	11.000	m	1.46	0.09
Sea state Bias	-0.500	0.000	m	1.39	0.22
Backscatter coefficient	7.000	30.000	dB	1.44	0.23
Ocean tide height	-5.000	5.000	m	0.01	0.15
Earth tide	-1.000	1.000	m	0.00	0.00
Pole tide	-15.000	15.000	m	0.00	0.00
TMR and ECMWF tropospheric differences	-0.200	0.200	m	NaN	0.35
Spline fitting					0.01

The following three maps are complementary: they show respectively the removed, the selected measurements and the percentage of selected measurements in the editing procedure.


Valid data
TOPEX/Poseidon Cycle 408 (11/10/2003 / 21/10/2003)



Edited measurements
TOPEX Cycle 408 (11/10/2003 / 21/10/2003)




Percentage of valid data relative to the nominal pass
TOPEX/Poseidon Cycle 408 (11/10/2003 / 21/10/2003)




3.5 Ionospheric correction

TOPEX/Poseidon Cycle 408 (11/10/2003 – 21/10/2003), ascending passes



TOPEX/Poseidon Cycle 408 (11/10/2003 – 21/10/2003), descending passes



minimum number of samples per box 1
 maximum number of samples per box 637



Statistics Y-X

mean = -0.01489
 rms = 0.02409
 std = 0.01894


Order 1 fit polynom

$y = a x + b$
 $a = 0.79496425$
 $b = 0.00281840$


Legend

— Order 1 fit polynom
 - - - Bisectrix

TOPEX/Poseidon Cycle 408 (11/10/2003 – 21/10/2003), ascending passes



minimum number of samples per box 1
 maximum number of samples per box 2401



Statistics Y-X

mean = -0.01344
 rms = 0.01744
 std = 0.01111


Order 1 fit polynom

$y = a x + b$
 $a = 0.41573662$
 $b = 0.00433876$


Legend

— Order 1 fit polynom
 — Bisectrix

TOPEX/Poseidon Cycle 408 (11/10/2003 – 21/10/2003), descending passes



minimum number of samples per box 1
 maximum number of samples per box 422



Statistics Y-X

mean = -0.01102
 rms = 0.02669
 std = 0.02431


Order 1 fit polynom

$y = a x + b$
 $a = 0.66831321$
 $b = 0.01776700$


Legend

— Order 1 fit polynom
 - - - Bisectrix

TOPEX/Poseidon Cycle 408 (11/10/2003 – 21/10/2003), ascending passes



minimum number of samples per box 1
 maximum number of samples per box 6065



Statistics Y-X

mean = -0.00090
 rms = 0.00724
 std = 0.00718


Order 1 fit polynom

$y = a x + b$
 $a = 0.72681528$
 $b = 0.00399030$


Legend

— Order 1 fit polynom
 — Bisectrix


TOPEX/Poseidon Cycle 408 (11/10/2003 – 21/10/2003), descending passes



3.6 Wet tropospheric correction



3.7 Crossover statistics



T1T1 – CROSSOVER STATISTICS

Without orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS


Type de points de croisement: T1T1
 Zone géographique (deg): -90 / 90 , 0 / 360
 Seuil sur les écarts d'analyse 0.00 (moy) 30.00 (seuil)
 Selection(s) sur les champs :
 CL Arc 1 :=INTERP_SPLN
 CL Arc 2 :=INTERP_SPLN
 Seuil Min +: 0.0000000
 Seuil Max : 0.0000000

Selection(s) sur les écarts :
 Aucune

RESULTATS STATISTIQUES

Valeur minimale : -29.8700
 Valeur maximale : 29.6600
 Différence Max – Min: 59.5300
 Nombre de points lus: 4785
 Nombre de points sélectionnés: 4670
 Moyenne : 0.772092
 Écart-type : 6.60671
 Moyenne Quadratique : 6.65167

CLS Space Oceanography Division



T1T1 – CROSSOVER STATISTICS

With orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS


Type de points de croisement: T1T1
 Zone géographique (deg): -90 / 90 , 0 / 360
 Seuil sur les écarts d'analyse 0.00 (moy) 30.00 (seuil)
 Selection(s) sur les champs :
 CL Arc 1 :=INTERP_SPLN
 CL Arc 2 :=INTERP_SPLN
 Seuil Min +: 0.0000000
 Seuil Max : 0.0000000

Selection(s) sur les écarts :
 Aucune

RESULTATS STATISTIQUES

Valeur minimale : -28.2500
 Valeur maximale : 29.9500
 Différence Max – Min: 58.2000
 Nombre de points lus: 4785
 Nombre de points sélectionnés: 4655
 Moyenne : 0.0157940
 Écart-type : 5.86668
 Moyenne Quadratique : 5.86670

CLS Space Oceanography Division



T1T1 – CROSSOVER STATISTICS
SSH, BATHY < -1000 m, VAR_OCE < 20 cm, LAT [-50°, +50]
SSH = Corrected sea surface height before orbit error


RAPPEL DES SELECTIONS

Type de points de croisement: T1T1
 Zone géographique (deg): -50 / 50 , 0 / 360
 Seuil sur les écarts d'analyse : aucun
 Selection(s) sur les champs :
 CL Arc 1 : =BATHY
 CL Arc 2 : =BATHY
 Seuil Min : aucun
 Seuil Max : -100000.00
 CL Arc 1 : =VAR_OCE
 CL Arc 2 : =VAR_OCE
 Seuil Min : aucun
 Seuil Max : 20.000000
 [...]
 Selection(s) sur les écarts :
 Aucune

RESULTATS STATISTIQUES

Valeur minimale : -27.6200
 Valeur maximale : 32.7200
 Différence Max – Min: 60.3400
 Nombre de points lus: 2637
 Nombre de points selectionnes: 2413
 Moyenne : 0.552333
 Ecart-type : 5.44728
 Moyenne Quadratique : 5.47521


CLS Space Oceanography Division




3.8 SSH variability

3.8.1 Sea Level Anomaly


TOPEX/Poseidon, cycle 408
Period : 11/10/2003 – 21/10/2003



TOPEX/Poseidon, cycle 408
Period : 11/10/2003 – 21/10/2003




TOPEX/Poseidon, cycle 408
Period : 11/10/2003 – 21/10/2003




3.8.2 Comparison to a precise Mean Sea Surface

The CLS (2001) MSS model is used as a reference to compute SLA. The two following maps respectively show the map of Topex SLA relative to the MSS and differences higher than a 30 cm threshold (after centering the data). The latter figure shows that higher differences are located in high ocean variability areas, as expected.

TOPEX/Poseidon, cycle 408
Period : 11/10/2003 – 21/10/2003




(SSH - MSS) differences greater than 0.3 m
TOPEX/Poseidon Cycle 408 (11/10/2003 / 21/10/2003)



3.9 Dynamic topography


TOPEX/Poseidon, cycle 408
Period : 11/10/2003 – 21/10/2003



3.10 Wind and wave maps

These two figures show wind and wave estimations derived from 10 days of altimeter measurements.

TOPEX/Poseidon, cycle 408
Period : 11/10/2003 – 21/10/2003



TOPEX/Poseidon, cycle 408
Period : 11/10/2003 – 21/10/2003

