


Coherent Lagrangian Pathways from the Surface Ocean to Interior




Ananda Pascual (IMEDEA, Spain) on behalf of the CALYPSO team

CALYPSO objectives

Unravel the three-dimensional coherent pathways by which water carrying tracers and drifting objects is transported from the surface ocean to depths below the mixed layer.






Ruiz et al. 2009

D'Asaro et al. 2018

Technical approach

- Focus on South Western Mediterranean
 - Strong meandering current and front
 - Deep subduction previously determined
- Experiments in July 2017, May/June 2018, March 2019
- Experiment planned for 2020/2021
- Modeling multi-scale approach
- Lagrangian perspective /coherent structure detection

WMOP - ROMS









CALYPSO meeting – Pollensa (Mallorca), 6-7 June 2019



CALYPSO meeting – Pollensa (Mallorca), 6-7 June 2019

Goals of the meeting:


- 1. Present preliminary results
- 2. Make plans for publication
- 3. Plan for 2020/2021 field work

- Presentations + working groups discussions and reports
- Synergies with other programs (SWOT Adopt a Cross-Over) and groups (MIO, SHOM)

E. D'Asaro (UW), A. Mahadevan (WHOI)


Chief Scientists

RV Pourquoi Pas? March 28 – April 11, 2019



S. Ruiz (IMEDEA)


Satellite context



S. Ruiz (IMEDEA)

Satellite context (altimetry)

2019




2018

E. D'Asaro (UW), A. Mahadevan (WHOI)


RV Pourquoi Pas?





LEG 1 : uCTD, ADCP, Microstructure, SVP & CODE Drifters, ARGO, CTD: flow cytometry, DNA








LEG 2 : u/ecoCTD, ADCP, profling floats, CTD: cytometry, DNA, Lagrangian floats, V-Wing, WireWalker, Drifters: SVP,CODE, CARTHE, WHOI

uCTD


E. Cutolo (IMEDEA)











Vertical Velocity from QG at 50m

3.4°W

3.2°W

3°W


2.8°W

2.6°W

0.00


3.8°W

3.6°W




uCTD


E. Cutolo (IMEDEA)



Moorings 2019


- Each measures temperature and salinity with 7 instruments in approximately upper 500 m.
- Real-time data delivery
- Geostrophic currents and transport. Upper 500 m, relative to 500 m.
- Transport is ± 1.5 Sv between pairs of moorings
- Signal reverses over duration of 6 days





Underwater glider surveys 2019

D. Rudnick (SIO)



20 March - 20 May 2019

400 glider-days


9300 km

2400 dives

Surface to 700 m

> 60 sections

Glider sections 2019




Southern section

Strong confluent front

Surface layer deepens to south


Suggestion of across-front convergence

P.-M.Poulain (CMRE), T. Ozgokmen (RSMAS), L. Centurioni (SIO)



Kinematic properties from drifters

Dataset from CALYPSO 2018 cruise – Least Square Method (Molinari et al. 1975; Essink et al. 2019)



Bin plots (scales 10-40 km, hourly estimations):

Drifters show coherent signals of convergence (order 0.5 f)

Vorticity values around 0.5 f.

CALYPSO 2018 E. D'Asaro (UW) Submesoscale front with shallow subduction Ship Survey and Mapped Potential Density Potential Density uCTD Section 152.4 0 a d 5 152.3 10 0 hours Pressure yearday 152.2 20 152.1 Float 25 Deploy km/ 10 km 152 30 -2.3 -2.28 -2.26 -2.24 -2.22 -2.2 -2.3 -2.28 -2.26 -2.24 -2.22 -2.2 longitude longitude Float Depth and Potential Density W mm/s 0 F b Color is vertical velocity 5 20 O(10 mm/s) 10 10 Depth (m) 19 hours more Pressure ADCP ~ 10 m on Float 15 weak events -FLOAT <3 m deep

10

12

18/06/01 05:12

20

25

30

152.1

152.15

1 hour

yearday

152.25

152.3

152.35

152.2



10

-20

30

05:36

05:24

~45 minutes

06:00

06:12

06:24


06:36


05:48

Time

E. D'Asaro (UW)

Simple Cartoon






We measure energetic upward and downward vertical motions near the surface with superinertial frequencies and submesoscale spatial scales.

Coherent structures?

Ruiz et al. 2009

This is consistent with models




Spectrum is roughly white Small scales dominate energy

These are "invisible" in SSH

McWilliams, Gula, Molemaker 2019

Kinematic properties from drifters

In the area of the float subduction – encouraging preliminary results



 Surface drifters show signals of convergence O(f)

• Vorticity values around 2 f.

PDF (scales <10km, 20 min estimations)

Summary and future plans

- Vertical velocity is complicated !
- Need of multiple 'toys'
- Further analysis (observations and modeling). Tighter coupling with forecasting models, data assimilation
- Potential synergies with SWOT
- Experimental plan 2020-2021 (drift-air, more floats, AUVs,...)
- Working groups (modeling, lagrangian, sampling strategy, location, biogeochemistry,..)
- Next meeting in 6 months

